

## What Are Prebiotics and Why Should You Care? (Part 1)

Peter Swann, MD, FAAFP, FACOEM

In previous articles, I spoke about the different kinds of fiber and their effects, and the potential risks of taking probiotics without also consuming prebiotic soluble fiber (PSF) in foods and/or supplements [see [August](#) & [October](#) 2016 issues]. That naturally raises the question of what PSF is and what scientific evidence supports the health effects of consuming it.

The concept of prebiotics was originally proposed by Gibson and Roberfroid in 1995, and defined as follows: "A non-digestible food ingredient that beneficially affects the host by selectively stimulating the growth and/or activity of one or a limited number of bacteria in the colon, and thus improves host health."<sup>1</sup>

Over the ensuing years, as the science progressed, the definition of prebiotics matured, with the most recent definition from the 6th Meeting of the International Scientific Association of Probiotics and Prebiotics stated as, "A dietary prebiotic is a selectively fermented ingredient that results in specific changes, in the composition and/or activity of the gastrointestinal microbiota, thus conferring benefit(s) upon host health."<sup>2</sup>

### Health Benefits

Many studies show positive health effects from prebiotic fiber supplementation, including the following:

- Improvements in the number and metabolic activity of healthful bacteria, primarily bifidobacteria and lactobacillus species
- Improvements in immune function and regulation
- Improvements in gastrointestinal disorders, including irritable bowel syndrome (IBS), inflammatory bowel disease (IBD) and decreased risk of colorectal cancer
- Improvement in mineral absorption, including calcium and magnesium, and increased bone density
- Decreased appetite and improved weight management in overweight subjects
- Improved glucose and lipid homeostasis

Let's take a stroll through the literature and explore the evidence for the first three above-mentioned health benefits, with the balance covered in a future article.

### Improvements in Number and Metabolic Activity of Healthful Bacteria

All health benefits start with "the prebiotic effect" - changes in the composition and/or metabolic activity of certain bacteria that have been shown to positively influence the health of the host. Beneficial effects can be related to their metabolism (i.e., fermentation profiles and end products), capacity for producing vitamins, antioxidants (reduction equivalents), bacteriocins against potentially harmful competitors, exchange of molecular signals between the different genera / species, but also with the eukaryotic epithelial cells.<sup>3</sup> The mechanism for those changes was shown

diagrammatically in [an article](#) in *Nature Biotechnology*, March 10, 2014.<sup>4</sup>

All types of prebiotic soluble fiber show certain characteristics:<sup>5</sup> resistance to gastric acidity, hydrolysis by mammalian digestive enzymes and GI absorption; fermentation by intestinal microflora; and selective stimulation of the growth and/or activity of one or a limited number of intestinal bacteria beneficially associated with health and well-being.

Many animal and human studies show that supplementation with a variety of types of dietary soluble fibers results in increases in bifidobacteria, lactobacillus or both.<sup>6-14</sup> Those and other healthful bacteria improve host health through a variety of metabolic intermediaries. They ferment dietary oligosaccharide soluble fiber, their primary food substrate, increasing in number and metabolic activity, yielding short-chain fatty acids (SCFA) butyrate, propionate and acetate, vitamins, antioxidants, and bacteriocins - narrow-spectrum natural antibiotics that help keep their competition at bay while simultaneously inhibiting some human colonic pathogens.

Butyrate serves as the primary food substrate for colonic epithelial cells (enterocytes), helping to keep them healthy and thus supporting the colonic epithelium's barrier function; and regulates enterocyte cell growth and differentiation - factors that may contribute to butyrate's role in reducing the risk of colon cancer.<sup>15</sup> Acetate and propionate, however, are mostly absorbed and transported to the liver, where they favorably affect glucose metabolism.<sup>16</sup> Propionate also appears to inhibit cholesterol synthesis and regulate adipose tissue deposition.<sup>17-18</sup>

#### Improvements in Immune Function and Regulation

The immune system is comprised of multiple, functionally different cell types, yielding a spectrum of immunomodulating molecules.<sup>19</sup> Although no single immune marker accurately reflects overall immune function, looking at many markers in different situations can paint a picture of immune function and modulation in both hyper-immune conditions (e.g., autoimmune disorders) and hypo-immune ones (e.g., infection.)

The interactive coexistence of the microbiota and the immune system is particularly clear in the colon, where gut-associated lymphoid tissue (GALT) has evolved to provide both defense against pathogens and tolerance of dietary and self-antigens. Colonic enterocytes are key intermediaries that convey signals from the intestinal lumen to the GALT.<sup>20-21</sup>

Non-breast-fed infants receiving prebiotic supplementation show a significantly higher concentration of fecal secretory IgA (sIgA) antibodies,<sup>22-23</sup> while in elderly adults, there is increased natural killer (NK) cell activity, increased production of anti-inflammatory cytokine IL-10, and decreased production of pro-inflammatory cytokines  $1\beta$ , IL-6 and TNF $\alpha$ . In infants with a high risk of allergies, prebiotic supplementation reduces plasma levels of total IgE, and IgG1,2 and 3, and lowers the incidence of atopic dermatitis.<sup>24</sup>

Experimental data from prebiotic supplementation animal studies show increased sIgA in the intestinal lumen, increased B cell numbers in Peyer's patches, enhanced production of anti-inflammatory IL-10, and decreased production of pro-inflammatory cytokines.<sup>25-26</sup> Additionally, in a review article, Lomax notes that many animal and human studies suggest some aspects of innate and adaptive immunity of the gut and systemic immune systems are positively modified by prebiotic supplementation, and a number have shown benefits in infectious outcomes.<sup>27</sup>

This body of work, along with other studies, taken together, strongly suggests prebiotic supplementation results in immunomodulation, improving the body's response to certain infections and improving inflammatory conditions.

#### Improvements in GI Disorders and Decreased Risk of Colorectal Cancer

IBS is a functional bowel condition of unknown etiology. Patients experience chronic, recurring abdominal pain or discomfort coinciding with diarrhea, constipation or both. In subjects with IBS, prebiotic supplementation has been shown to significantly reduce the incidence and severity of symptoms while increasing ratings of quality of life.<sup>28-30</sup>

IBD, on the other hand, is a chronic, relapsing disorder characterized by intestinal inflammation, ulceration and stricturing resulting from alterations in the balance of proinflammatory and immunoregulatory cytokines within the mucosal immune system. Symptoms include diarrhea, fecal urgency and incontinence, severe abdominal pain and rectal bleeding, and can have a profound negative impact on nutritional status and quality of life.

Studies on prebiotic supplementation in patients with ulcerative colitis (IC) have shown significant reductions in inflammation, with increased mucosal bifidobacteria and decreased pro-inflammatory TNF $\alpha$  and IL-10 $\alpha$ ;<sup>31-32</sup> and similar changes in [Crohn's disease](#) patients accompanied by a significant reduction in disease activity, with 40 percent of the treated patients entering disease remission.<sup>33</sup>

Evidence from a wide variety of sources implicates alterations in the colonic microbiota in the etiology of colon cancer.<sup>34-35</sup> Therefore, it follows that improving the number and activity of healthful gut microbes may interfere with the process of carcinogenesis. Animal studies on prebiotic supplementation have shown decreased pre-cancerous lesions<sup>36-39</sup> and decreased tumor incidence.<sup>40-42</sup> For human intervention trials, cancer is an impractical end point in terms of number of subjects, study duration, cost and ethical considerations. Human studies, therefore, rely on changes in biomarkers of colon cancer such as DNA damage and cell proliferation in colonic mucosa.<sup>43</sup>

A randomized, double-blind, trial of prebiotic supplementation in patients with resected polyps or colon cancer showed favorable changes in multiple colon cancer biomarkers.<sup>44</sup> This body of work, along with other studies, taken together, strongly suggest prebiotic supplementation helps prevent or ameliorate multiple colonic conditions, including IBS, IBD and colorectal cancer.

#### References

1. Gibson GR, Roberfroid M. Dietary modulation of the human colonic microbiota: introducing the concept of prebiotics. *J Nutrition*, July 1995;125(6):1401-12.
2. Gibson GR, et al. Dietary prebiotics: current status and new definition. *Food Sci Tech Bulletin: Functional Foods*, May 2010;7(1):1-19.
3. Roberfroid M, et al. Prebiotic effects: metabolic and health benefits. *British J Nutrition*, August 2010;104(Suppl 2):S1-63.
4. Brüssow H, Parkinson SJ. "You Are What You Eat." *Nature Biotech*, March 2014;32(3):243-45.
5. Gibson GR, Roberfroid M. Dietary modulation of the human colonic microbiota: introducing the concept of prebiotics. *J Nutrition*, June 1995;125(6):1401-12.
6. Kruse H-P, et al. Effects of inulin on faecal bifidobacteria in human subjects. *British J Nutrition*, 1999;82(5):375-82.

7. Tuohy KM, et al. The prebiotic effects of biscuits containing partially hydrolysed guar gum and fructo-oligosaccharides - a human volunteer study. *British J Nutrition*, September 2001;86(3):341.
8. Menne E, Guggenbuhl N, Roberfroid M. "Fn-type chicory inulin hydrolysate has a prebiotic effect in humans. *J Nutrition*, May 2000;130(5):1197-99.
9. Moro G, et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatric Gastroenterology & Nutrition*, 2002;34(3):291-95.
10. Bouhnik Y, et al. The capacity of short-chain fructo-oligosaccharides to stimulate faecal bifidobacteria: a dose-response relationship study in healthy humans. *Nutrition J*, January 2006:8.
11. Lee PR, et al. Dietary treatments for epilepsy: management guidelines for the general practitioner. *Epilepsy & Behavior*, June 2011;21(2):115-21.
12. Yen C-H, et al. Long-term supplementation of isomalto-oligosaccharides improved colonic microflora profile, bowel function, and blood cholesterol levels in constipated elderly people — a placebo-controlled, diet-controlled trial. *Nutrition*, April 2011;27(4):445-50.
13. Thitaram SN, et al. Isomaltooligosaccharide increases cecal bifidobacterium population in young broiler chickens. *Poultry Science*, July 2005;84(7):998-1003.
14. Day D, Chung C-H. Isomaltooligosaccharides from leuconostoc as neutraceuticals: <https://patents.google.com/patent/US7291607B2/en>.
15. Roberfroid M, et al., *Op Cit* (Reference #3).
16. Brüssow H, et al., *Op Cit* (Reference #4).
17. Reshef L, et al. Effect of propionate on lipogenesis in adipose tissue. *J Lipid Research*, November 1967;8(6):682-87.
18. Xiong Y, et al. Short-chain fatty acids stimulate leptin production in adipocytes through the G protein-coupled receptor GPR41. *Proceed National Academy Sci*, January 2004;101(4):1045-50.
19. Murphy K, Weaver C. *Janeway's Immunobiology, 9th Edition*. New York: Garland Publishing, 2016.
20. Sanderson IR. Dietary modulation of GALT. *J Nutrition*, November 2007;137(11 Suppl):2557S-2562S.
21. Artis D. Epithelial-cell recognition of commensal bacteria and maintenance of immune homeostasis in the gut. *Nature Reviews Immunology*, June 2008;8(6):411-20.
22. Petra AM, Scholtens J, et al. Fecal secretory immunoglobulin A is increased in healthy infants who receive a formula with short-chain galacto-oligosaccharides and long-chain fructo-oligosaccharides. *J Nutrition*, June 2008;138(6):1141-1147.  
<http://jn.nutrition.org/content/138/6/1141.full>
23. Bakker-Zierikzee AM, et al. Faecal SIgA secretion in infants fed on pre- or probiotic infant formula. *Pediatric Allergy and Immunology*, March 2006;17(2):134-40.
24. Moro G, et al. A mixture of prebiotic oligosaccharides reduces the incidence of atopic dermatitis during the first six months of age. *Archives of Disease in Childhood*, October 2006;91(10):814-19.
25. Seifert S, Watzl B. Inulin and oligofructose: review of experimental data on immune modulation. *J Nutrition*, November 2007;137(11 Suppl):2563S-2567S.
26. Watzl B, et al. Inulin, oligofructose and immunomodulation. *British J Nutrition*, April 2005;93(S1):S49.
27. Lomax AR, Calder PC. Prebiotics, immune function, infection and inflammation: a review of the evidence. *British J Nutrition*, March 2009;101(5):633-58.
28. Paineau D, et al. The effects of regular consumption of short-chain fructo-oligosaccharides on digestive comfort of subjects with minor functional bowel disorders. *British J Nutrition*, February 2008;99(2):311-18.
29. Silk DBA, et al. Clinical trial: the effects of a trans-galactooligosaccharide prebiotic on faecal microbiota and symptoms in irritable bowel syndrome. *Alimentary Pharmacology & Therapeutics*, March 2009;29(5):508-18.

30. Buddington KK, et al. Dietary oligofructose and inulin protect mice from enteric and systemic pathogens and tumor inducers. *J Nutrition*, March 2002;132(3):472-77.
31. Furrie E, et al. Symbiotic therapy (bifidobacterium longum/synergy 1) initiates resolution of inflammation in patients with active ulcerative colitis: a randomised controlled pilot trial. *Gut*, February 2005;54(2):242-49.
32. Casellas F, et al. Oral oligofructose-enriched inulin supplementation in acute ulcerative colitis is well tolerated and associated with lowered faecal calprotectin. *Alimentary Pharmacology & Therapeutics*, 2007;25(9):1061-67.
33. Lindsay JO, et al. Clinical, microbiological, and immunological effects of fructo-oligosaccharide in patients with Crohn's disease. *Gut*, March 2006;55(3):348-55.
34. Rowland IR. The role of the gastrointestinal microbiota in colorectal cancer. *Current Pharmaceutical Design*, 2009;15(13):1524-27.
35. Cipe G, et al. Relationship between intestinal microbiota and colorectal cancer. *World J Gastrointestinal Oncology*, October 2015;7(10):233-40.
36. Verghese M, et al. Dietary inulin suppresses azoxymethane-induced preneoplastic aberrant crypt foci in mature fisher 344 rats. *J Nutrition*, September 2002;132(9):2804-8.
37. Reddy BS, et al. Effect of dietary oligofructose and inulin on colonic preneoplastic aberrant crypt foci inhibition. *Carcinogenesis*, July 1997;18(7):1371-74.
38. Poulsen M, et al. Different effects of short- and long-chained fructans on large intestinal physiology and carcinogen-induced aberrant crypt foci in rats. *Nutrition and Cancer*, 2002;42(2):194-205.
39. Buddington D, et al., *Op Cit* (Reference #30).
40. Jacobsen H, et al. Carbohydrate digestibility predicts colon carcinogenesis in azoxymethane-treated rats. *Nutrition and Cancer*, July 2006;55(2):163-70.
41. Caderni G, et al. Identification of mucin-depleted foci in the unsectioned colon of azoxymethane-treated rats: correlation with carcinogenesis. *Cancer Research*, May 2003;63(10):2388-92.
42. Pierre F, et al. Short-Chain Fructo-Oligosaccharides Reduce the Occurrence of Colon Tumors and Develop Gut-Associated Lymphoid Tissue in Min Mice., *Cancer Research*, January 1997;57(2):225-28.
43. Roberfroid M, et al., *Op Cit* (Reference #3).
44. Femia AP, et al. Antitumorigenic activity of the prebiotic inulin enriched with oligofructose in combination with the probiotics lactobacillus rhamnosus and bifidobacterium lactis on azoxymethane-induced colon carcinogenesis in rats. *Carcinogenesis*, November 2002;23(11):1953-60.

JANUARY 2017