# Dynamic Chiropractic



EVIDENCE / RESEARCH / SCIENCE

## Monochromatic Infrared Energy for Diabetic Neuropathy?

Ronald Feise, DC

## WHAT YOU NEED TO KNOW

- Theoretically, devices delivering this therapy are used to increase circulation, reduce pain and improve neuropathology symptoms.
- Currently, five randomized clinical trials with placebo controls have investigated the effects of monochromatic infrared energy. Unfortunately, four found no clinical benefit compared with placebo and the fifth found conflicting results.
- A conservative package of care using six main strategies (three lifestyle interventions, one topical intervention and two oral interventions) is recommended instead.

As clinical research director for numerous chiropractic clinics, I am asked to consult in some very interesting cases. A recent patient had been experiencing chronic painful diabetic neuropathy for over three years and had been treated with monochromatic infrared energy for six months. None of her symptoms had improved. In fact, the patient reported that her feet were more painful and her paresthesia had worsened since she began care. The patient wanted our opinion on the treatment she had received and, more importantly, what treatments we would recommend.

#### Diabetic Neuropathy: The Stats

The natural history of diabetic neuropathy is progressive and irreversible loss of sensibility in the feet, leading to ulceration and/or amputation in 15% of patients. The prevalence of neuropathy is

more than 50% in those who have been diabetic for 20 years.<sup>1</sup> Worldwide, more than 420 million patients currently have diabetes.

A five-year follow-up study of community patients with chronic painful diabetic neuropathy demonstrated that 23% of patients can experience complete remission of neuropathic symptoms,

but most patients will continue to experience pain and paresthesia.<sup>5</sup>

### **Clinical Assessment**

Clinical history alone is an insufficient screen to for diabetes-related peripheral neuropathy,

because half of patients with diabetes-related peripheral neuropathy are asymptomatic.<sup>14</sup> Once there is a clinical suspicion of diabetic neuropathy, a thorough clinical assessment must exclude other causes of neuropathy, and should involve a comprehensive history and examination including: 10 g monofilament, 128 Hz tuning fork, Ipswich touch test, temperature perception, vibration perception threshold, pinprick, proprioception and ankle reflexes. You should use a measure to help monitor patient progress.

The Neuropathic Pain Symptom Inventory has been designed specifically to monitor effects of therapy on neuropathic pain.<sup>2</sup>

Monochromatic Infrared Energy: Poor Research Support

Over the past 20 years, monochromatic infrared energy has gained popularity in health care. Theoretically, devices delivering this therapy are used to increase circulation, reduce pain and improve neuropathology symptoms.

The initial evidence regarding the benefit of monochromatic infrared energy has been limited, because early studies used poor research methods that produced clouded findings. Early uncontrolled studies failed to account for placebo effects or the natural history.

It is important to note that the newer studies with a placebo arm have demonstrated a strong placebo effect for monochromatic infrared energy. The placebo effect and natural history explain why early studies (case reports, uncontrolled studies) and clinical observations mistakenly found evidence of effectiveness for monochromatic infrared energy.

Currently, five randomized clinical trials with placebo controls have investigated the effects of monochromatic infrared energy.<sup>4, 8-11</sup> Unfortunately, four of the five studies found no clinical benefit compared with placebo controls, and the fifth study found conflicting results.

### A Better (Conservative) Protocol

We could not endorse monochromatic infrared energy for this patient with chronic, painful diabetic neuropathy. Instead, we recommended a conservative package of care using six main strategies (three lifestyle interventions, one topical intervention and two oral interventions), strict control of diabetes, smoking cessation, exercise, topical capsaicin, acetyl-L-carnitine and  $\alpha$ -lipoic acid.

#### ASSESSMENT OF SHOCKWAVE, ELECTRO-STIMULATION, AND LOW-LEVEL LASER IN PATIENTS WITH DIABETIC NEUROPATHY

Although shockwave, electrostimulation and low-level laser are effective for many musculoskeletal disorders, evidence is lacking for their use in patients with diabetic neuropathy.

This assessment excluded studies with fatal research fl aws [i.e., absence of randomized clinical trial design with either active or placebo control group comparison, small samples (20%)]. Serious fl aws erode the fi ndings of a study; thus, fi ndings of fl awed studies should be demoted when making clinical decisions for our patients.

Shockwave: Animal studies using shockwave demonstrate a benefi t. However, there are no human trials demonstrating a benefi t. $^{3-6}$ 

Electrostimulation: Most electrostimulation studies suffer from poor research quality and yield mixed results (some demonstrate improved pain and quality of life, and others do not). Two quality randomized clinical trials demonstrated confl icting outcomes. Bosi, et al., showed improvements in pain reduction and quality of life; but Khalil, et al., found that electrostimulation was not better than a placebo for changes in sensory nerve function.<sup>2,4</sup>

Low-Level Laser: Most low-level laser studies suffer from poor research quality. Two quality randomized clinical trials examined laser and showed inconsistent results. Swislocki, et al., showed mixed results: there were improvements in qualityof-life items, but not in pain intensity.<sup>5</sup> However, Bashiri demonstrated that laser had a beneficial effect on both pain and quality-of-life patient outcomes.<sup>1</sup>

Note: Sidebar citations are separately numbered from in-text citations (both reference lists accompany the web version of this article).

Strict control of diabetes has robust evidence in reducing the risk of developing diabetic peripheral neuropathy. The mainstay of modern management is glycemic control to manage and prevent diabetes complications.<sup>17</sup> There is a positive association between smoking and the prevalence and incidence of diabetic peripheral neuropathy.<sup>3</sup> It has been demonstrated that smoking cessation improves neuropathy symptoms.<sup>15</sup>

A systematic review shows that a combination of aerobic and sensorimotor training also is beneficial in improving symptoms.<sup>13</sup> The aerobic exercise can be of moderate-intensity (40-70%

max heart rate), 30 minutes per day, performed three to six times per week.

Overall, randomized clinical trials support the effects of topical capsaicin applied to the feet in

improving pain and other neuropathy symptoms.<sup>7</sup> The studies used four applications/day for eight weeks, which can be extended. Capsaicin cream is well-tolerated, but local skin reactions were common and, if severe, you prompt discontinuation of treatment.

Multiple randomized clinical trials found that acetyl-L-carnitine at doses greater than 1,500mg/day

reduced pain more than placebo.<sup>6, 12</sup> Alpha-lipoic acid was well-tolerated in randomized clinical

trials and was associated with improvement of neuropathic impairments and deficits.<sup>18-19</sup> A Mayo Clinic review recommends the following dosing: acetyl-L-carnitine 1,000 mg, 3x/day; and  $\alpha$ -lipoic acid 600 mg/day.<sup>16</sup>

#### References

- 1. Aszmann O, Tassler PL, Dellon AL. Changing the natural history of diabetic neuropathy: incidence of ulcer/amputation in the contralateral limb of patients with a unilateral nerve decompression procedure. *Ann Plast Surg*, 2004;53:517-22.
- 2. Bouhassira D, Attal N, Fermanian J, et al. Devel-opment and validation of the Neuropathic Pain Symptom Inventory. *Pain*, 2004;108:248-57.
- 3. Clair C, Cohen MJ, Eichler F, et al. The effect of cigarette smoking on diabetic peripheral neuropathy: a systematic review and meta-analysis. *J Gen Intern Med*, 2015;30:1193-203.
- 4. Clifft JK, Kasser RJ, Newton TS, Bush AJ. The effect of monochromatic infrared energy on sensation in patients with diabetic peripheral neuropathy: a double-blind, placebo-controlled study. *Diabetes Care*, 2005;28:2896-900.
- 5. Daousi C, Benbow SJ, Woodward A, MacFarlane IA. The natural history of chronic painful peripheral neuropathy in a community diabetes population. *Diabet Med*, 2006;23:1021-4.
- 6. De Grandis D, Minardi C. Acetyl-L-carnitine (levacecarnine) in the treatment of diabetic neuropathy. A long-term, randomised, double-blind, placebo-controlled study. *Drugs R&D*, 2002;3:223-31.
- 7. Dludla PV, Nkambule BB, Cirilli I, et al. Capsaicin, its clinical significance in patients with painful diabetic neuropathy. *Biomed Pharmacother*, 2022;153:113439.
- 8. Franzen-Korzendorfer H, Blackinton M, Rone-Adams S, McCulloch J. The effect of monochromatic infrared energy on transcutaneous oxygen measurements and protective sensation: results of a controlled, double-blind, randomized clinical study. *Ostomy Wound Manage*, 2008;54:16-31.
- 9. Lavery LA, Murdoch DP, Williams J, Lavery DC. Does anodyne light therapy improve peripheral neuropathy in diabetes? A double-blind, sham-controlled, randomized trial to evaluate monochromatic infrared photoenergy. *Diabetes Care*, 2008;31:316-21.
- 10. Leonard DR, Farooqi MH, Myers S. Restoration of sensation, reduced pain, and improved balance in subjects with diabetic peripheral neuropathy: a double-blind, randomized, placebo-controlled study with monochromatic near-infrared treatment. *Diabetes Care*, 2004;27:168-72.
- 11. Nawfar SA, Yacob NB. Effects of monochromatic infrared energy therapy on diabetic feet with peripheral sensory neuropathy: a randomised controlled trial. *Singapore Med J*, 2011;52:669-72.
- 12. Sima AA, Calvani M, Mehra M, Amato A; Acetyl-L-Carnitine Study Group. Acetyl-L-carnitine improves pain, nerve regeneration, and vibratory perception in patients with chronic diabetic neuropathy: an analysis of two randomized placebo-controlled trials. *Diabetes Care*, 2005;28:89-94.
- 13. Streckmann F, Balke M, Cavaletti G, et al. Exercise and neuropathy: systematic review with meta-analysis. *Sports Med*, 2022;52:1043-1065.

- 14. Tesfaye S, Boulton AJ, Dyck PJ, et al. Diabetic neuropathies: update on defini- tions, diagnostic criteria, estimation of severity, and treatments. *Diabetes Care*, 2010;33:2285-93.
- 15. Voulgari C, Katsilambros N, Tentolouris N. Smoking cessation predicts amelioration of microalbuminuria in newly diagnosed type 2 diabetes mellitus: a 1-year prospective study. *Metabolism*, 2011;60:1456-64.
- 16. Watson JC, Dyck PJ. Peripheral neuropathy: a practical approach to diagnosis and symptom management. *Mayo Clin Proc*, 2015;90:940-51.
- 17. Yang H, Sloan G, Ye Y, et al. New perspective in diabetic neuropathy: from the periphery to the brain, a call for early detection, and precision medicine. *Front Endocrinol*, 2020;10:929.
- 18. Ziegler D, Ametov A, Barinov A, et al. Oral treatment with alpha-lipoic acid improves symptomatic diabetic polyneuropathy: the SYDNEY 2 trial. *Diabetes Care*, 2006;29:2365-70.
- Ziegler D, Low PA, Litchy WJ, et al. Efficacy and safety of antioxidant treatment with α-lipoic acid over 4 years in diabetic polyneuropathy: the NATHAN 1 trial. *Diabetes Care*, 2011;34:2054-60.

#### Sidebar References

- 1. Bashiri H. Evaluation of low level laser therapy in reducing diabetic polyneuropathy related pain and sensorimotor disorders. *Acta Med Iran*, 2013;51:543-7.
- 2. Bosi E, Conti M, Vermigli C, et al. Effectiveness of frequency-modulated electromagnetic neural stimulation in the treatment of painful diabetic neuropathy. *Diabetologia*, 2005;48:817-23.
- 3. Chen YL, Chen KH, Yin TC, et al. Extracorporeal shock wave therapy effectively prevented diabetic neuropathy. *Am J Transl Res*, 2015;7:2543-60.
- 4. Khalil Z, Ogrin R, Darzins PJ. The effect of sensory nerve stimulation on sensory nerve function in people with peripheral neuropathy associated with diabetes. *Neurol Res*, 2007;29:743-8.
- 5. Swislocki A, Orth M, Bales M, et al. A randomized clinical trial of the effectiveness of photon stimulation on pain, sensation, and quality of life in patients with diabetic peripheral neuropathy. *J Pain Symptom Manage*, 2010;39:88-99.
- 6. Zhou Y, Dai H, Long J, et al. Effects of extracorporeal shock waves on neuralgia in diabetic rats. *J Pain Res*, 2019;12:387-394.

SEPTEMBER 2023

©2024 Dynanamic Chiropractic<sup>™</sup> All Rights Reserved