

Chinese Skullcap: A Viable Adjunct to Cancer Treatment

James P. Meschino, DC, MS

One of the most respected and utilized herbs in traditional Chinese medicine (TCM) is Chinese skullcap (*scutellaria baicalensis*). Although less popular among Western health practitioners, the bioactive agents (e.g., baicalein flavonoid) in this herb have been the subject of intensive study with regard to their anticancer effects and other health benefits. In Asia, Chinese skullcap is incorporated into a number of combination herbal products that have been used in adjunctive cancer treatment. I recently had the opportunity to review the literature published worldwide on this potentially important herbal agent, and have summarized it below in a clinically relevant format. I believe you will find the data quite intriguing and of great clinical importance.

General Features

Chinese skullcap is a member of the mint family and grows in China and Russia.^{1,2} Its root is a rich source of over 35 flavonoids, giving it a yellow color - hence its traditional name of golden root or *huang qin*, the Chinese term for yellow gold (known as *ogon* in Japanese).^{3,4} One of the major flavonoids contained in the root of the plant is baicalin, a flavone glycoside (12percent-17percent of all skullcap root flavonoids), which yields the aglycone flavone baicalein, once hydrolyzed by gut bacteria.³ Baicalein is readily absorbed into the bloodstream, where it has been shown to exert a broad spectrum of important biological activities.⁵ Baicalein has been incorporated into a number of herbal combinations in TCM that treat a wide array of health conditions, including the widely used Asian herbal remedy, Sho-saiko-to.^{6,7} Currently, baicalein is being studied for its anti-cancer, anti-inflammatory, antiviral, antibacterial and anti-allergy effects.^{8,9,10,11} As explained below, its ability to halt the replication of various human cancer cell lines, via the inhibition of the 12 lipoxygenase enzyme system, is attracting a great deal of attention from the scientific community as a premiere agent in cancer research.

Clinical Application and Mechanism of Action: Anticancer Effects

Many experimental studies indicate that baicalein (5,6,7-trihydroxyflavone) prevents and inhibits cancer growth via a number of direct and indirect physiological actions:

a. Inhibition of 12-lipoxygenase enzyme and induction of apoptosis of cancer cells. Baicalein has been shown to inhibit the 12-lipoxygenase enzyme, which converts arachidonic acid into a specific leukotriene (eicosanoid) required for cancer cells to proliferate. Studies demonstrate that by inhibiting the enzyme, baicalein inhibits cancer cell proliferation and induces apoptosis (programmed cell death) of human gastric cancer cell lines.¹² Further, activation of arachidonic acid by the enzyme has been shown to be critical for metastasis of human prostate cancer cells.

Experimental evidence demonstrates that after intraprostatic injection of mice with human prostate cancer cells, the cells pretreated with baicalein failed to metastasize to the lung and failed

to express activity of the enzyme. This evidence suggests that in the presence of baicalein, various cancer cells can be prevented from multiplying and metastasizing to other tissues, and that a primary mechanism through which this occurs is via the inhibition of the enzyme.¹³ The inhibition of cancer cell proliferation also lends itself to increased apoptosis and a better opportunity for immune cells to destroy tumor cells.^{12,13}

Other studies testing baicalein as an antitumor agent have shown that it selectively induces apoptosis in human hepatoma cell lines (liver cancer cells) with minimal influence on noncancer cells.^{14,15,28,29} Its ability to induce apoptosis is under intensive investigation; many studies provide evidence that it exhibits an extraordinary ability to selectively encourage apoptosis of many different human cancer cells. This is likely one of the primary ways in which it has been a useful addition to herbal combinations used in cancer treatment.^{16,17,18}

b. Antiproliferative effects: Baicalein also has demonstrated anti-proliferative effects on human bladder cancer cell lines and a murine bladder cancer cell line, *in vitro*. In an *in vivo* experiment, mice were injected with bladder cancer cells with concurrent oral administration of a high-baicalein-yielding supplement in one group, or with no baicalein supplementation in the control group. All the control mice showed a progressive increase in tumor volume over the ensuing days of the study, whereas the mice treated with baicalein (*scutellaria*) showed a significant inhibition of tumor growth.¹⁹

In other experiments, baicalein has demonstrated antiproliferative effects on human pancreatic cancer cell lines and T lymphoid leukemia cells. The mechanism of action was shown to be through the reduction of protein tyrosine kinase activity and protein kinase C activity. Both of these enzyme activities are required for normal cellular proliferation to occur. Thus, in addition to inhibiting the 12-lipoxygenase enzyme pathway and inducing apoptosis of cancer cells, baicalein is also known to reduce cancer cell proliferation by directly or indirectly inhibiting specific enzymes (protein tyrosine kinase and protein kinase C) required for cellular division and proliferation.²⁸

c. Antioxidant effects: Baicalein is also a strong antioxidant and has been shown to protect DNA from undergoing cancerous mutations in challenge studies using potent carcinogens, such as benzo{a}pyrene, and toxins such as aflatoxin (AF)B.^{1,20,21}

d. Stimulation of DNA repair enzymes. *In vitro* evidence indicates baicalein stimulates recombination and repair of damaged DNA, supporting its traditional inclusion in cancer formulas, and suggesting possible use after sunburn and radiation damage.²²

e. Inhibition of the 5 alpha-reductase enzyme. Baicalein has been shown to inhibit the 5 alpha-reductase enzyme, which converts testosterone to dihydrotestosterone (DHT), strongly associated with the development of prostate enlargement (benign prostatic hyperplasia) and prostate cancer. As such, it is reported to be potentially useful for the prevention and/or treatment of androgen-dependent (testosterone-driven) disorders, including prostate enlargement and prostate cancer.²³

Human Studies Involving Prostate Cancer Patients

Human studies have primarily involved patients with advanced prostate cancer who were shown to be unresponsive to traditional medical drugs and other interventions. In a study performed by researchers from the University of California at San Francisco and Memorial Sloan-Kettering Cancer Center in New York, an herbal combination containing baicalein was administered orally to

metastatic prostate cancer patients (previously unresponsive to standard treatments). It was shown to demonstrate reversal and/or stabilization of their condition - with significantly improved survival, quality-of-life scores and other positive outcomes in almost all patients.

After a median treatment period of 57 weeks, all patients with androgen-dependent cancer had a decline of 80 percent or more in their prostate-specific antigen (PSA) levels, undetectable in 81 percent of the subjects. In 31 of 32 patients, testosterone levels declined to that seen in castrated men. Nine men with elevated prostatic acid phosphatase levels (a marker for cancer progression) experienced declines of more than 50 percent. Of two patients with positive bone scans, one experienced a complete disappearance of cancerous bone lesions, and the other showed improvement. One patient experienced a complete disappearance of a bladder mass, confirmed through a CT scan.

Of 35 patients with androgen-independent prostate cancer, 54 percent showed a drop in their PSA levels of 50 percent or more, and of 25 patients with metastasis to bone, two showed marked improvement; seven remained stable; 11 progressed; and five had no further bone scan follow-up, due to an increasing trend in their PSA levels.²⁴ Other studies involving men with known prostate cancer who were taking this herbal combination have demonstrated similar results. In a two-case report of hormone-refractory prostate cancer,²⁵ PSA levels dropped from 100ng/mL to 24, and from 386ng/mL to 114 within 16 months of supplementation. Studies following larger groups of men with prostate cancer for up to four years have also shown that this herbal formula significantly reduces PSA levels and improves survival and quality-of-life scores.^{26,27}

Researchers attribute much of the antitumor effects of this herbal combination to the presence of baicalein. According to researchers, it has demonstrated impressive anticancer effects against androgen-dependent and androgen-independent human prostate cancer cell lines, and many of its anticancer mechanisms have been uncovered in recent years, through in vitro and in vivo experiments.^{16-18,24-28} It appears to be of particular benefit in prostate cancer treatment, and in combination with other phytonutrients (plant-based nutrients) and micronutrients (vitamins and minerals), may help to prevent the development of clinically important prostate cancer, reducing prostate cancer incidence.²⁶ Animal studies indicate that baicalein may be effective in the treatment of other cancers as well, with experimental evidence supporting its potential use in stomach, pancreatic, bladder, liver^{12, 14, 15,19, 28} and breast cancer.²⁹⁻³¹

Hepatoma and Leukemia Patients

Based on its traditional use and the documentation of its antitumor effects against various human cancer cell lines, baicalein has been used recently by doctors in Asia as a complementary supplemental agent in the treatment of hepatomas and leukemia in human subjects.²⁹

Clinical Application and Mechanism of Action: Antiviral/Anti-HIV Effects

In experimental studies, baicalein has been shown to inhibit the activity of the HIV-1 integrase enzyme, essential to the life cycle of the human immunodeficiency virus (HIV). This enzyme is responsible for catalyzing the insertion of the viral genome into the host cell chromosome. It is an attractive target for the design of an HIV antiviral drug, because the integrase enzyme has no human counterpart.³² In a review of all herbal medicines frequently used as alternative medical therapies by HIV-positive and acquired immune deficiency syndrome (AIDS) patients, J.A. Wu, et al., emphasized that experimental data suggest that baicalein inhibits infectivity and replication of

HIV, and shows great promise as an important component to be included in herbal remedies used for HIV treatment.³³

Effects on Asthma and Allergy

Baicalein inhibits type I and II hypersensitivity reactions, confirming its traditional use in asthma^{34,35} and allergic dermatitis.³

Anti-Inflammatory Effects

Baicalein has shown impressive anti-inflammatory effects that appear to be mediated via repression of leukotriene B4 (inhibition of the 5-lipoxygenase enzyme), inhibition of interleukin-1B, and prostaglandin E2. Studies reveal it may be of benefit in the treatment of gingivitis (gum inflammation).^{29,36,37}

Effects on Alzheimer's and Dementia

In a rat model, baicalein has been shown to prevent the buildup of amyloid beta peptide (abeta), known to generate free-radical damage to brain cells and participate to a significant degree in the development and progression of Alzheimer's disease. Additionally, chronic inflammation occurs in Alzheimer's pathogenesis, and baicalein was shown to inhibit the 12-lipoxygenase enzyme responsible for brain inflammation, to a large degree. When tested against other known 12-lipoxygenase inhibitor agents, only baicalein was able to attenuate both nerve cell death (apoptosis) and the overexpression of abeta production. As such, it may help to prevent or forestall the development of Alzheimer's disease by reducing the buildup of abeta and inflammatory mediators, preventing brain cell death, which has been shown to be triggered by the accumulation of abeta in affected brain cells.³⁸

In studies using human neuroblastoma cells, the antioxidant properties of baicalein were shown to inhibit free-radical damage to these nerve cells induced by treatment with hydrogen peroxide (a powerful free-radical generating agent). The researchers argue that oxidative (free-radical) stress plays an important role in the development of neurodegenerative diseases (e.g., Alzheimer's disease, multiple sclerosis, Parkinson's disease) and that antioxidants such as baicalein and quercetin (also a flavonoid) may be useful bioactive agents in the prevention and/or management of these conditions, pending further studies. Both have shown impressive protective effects under experimental conditions.³⁹

Inhibition of Xanthine Oxidase (Gout)

Baicalein is one of few bioactive agents shown to inhibit the enzyme xanthine oxidase, responsible for the conversion of hypoxanthine to xanthine to uric acid, in the pathway for degradation of purines in the body. High levels of uric acid are a hallmark feature of gout and thus, baicalein supplementation may be an effective complementary intervention in the long-term management of this condition.⁴⁰ Allopurinol is a xanthine oxidase inhibitor, and a primary medication prescribed for the treatment of gout.

Dosage

Therapeutic purposes: The usual doses, for therapeutic purposes, range from 150-200 mg per day of Baicalein.⁶

General wellness: Consider 50-100mg per day.

Toxicity and Adverse Side Effects

Baicalein supplementation at therapeutic doses appears to be safe, with no reports of significant side-effects or toxicity. Baicalein or Chinese skullcap is not known to be contraindicated for any health conditions.^{41,42}

Drug-Nutrient Interaction

There are no known drug-nutrient interactions for baicalein or Chinese skullcap.⁴²

1. Bone K. Clinical Applications of Ayurvedic and Chinese herbs: Monographs for the Western herbal practitioner. Warwick, Australia 1996;p.75-9
2. Newall CA, Anderson LA, Phillipson JD. *Herbal Medicines: A Guide For Health-Care Professionals*. London; Pharmaceutical Press 1996;p.239-40
3. Tang W, Eisenbrand G. *Chinese Drugs of Plant origin: Chemistry, Pharmacology, and Use in Traditional and Modern Medicine*. New York; Springer-Verlag 1992.
4. Hsu HY. *Oriental Materia Medica: A Concise Guide*. Long Beach CA; Oriental Healing Arts Institute 1986.
5. Nishioka Y, Kyotani S, Miyamura M, Kusunose M. Influence of time of administration of Shosaiko-To extract granule on blood concentration of its active constituents. *Chem Pharm Bull* 1992; 40:p.1335-7.
6. Chen S, Ruan Q, Bedner E, Deptala A, Wang X, Hsieh TC, et al. Effects of the flavonoid baicalin and its metabolite baicalein on androgen receptor expression, cell cycle progression and apoptosis of prostate cancer cell lines. *Cell Prolif* 2001 Oct; vol 34(5):p.293-304.
7. Rossi M, Meyer R, Constantinou P, Caruso F, Castelbuono D, O'Brien M, et al. Molecular structure and activity toward DNA of baicalein, a flavone constituent of the Asian herbal medicine 'Sho-saiko-to. *J Nat Prod* 2001 Jan; vol 64(1):p.26-31.
8. Kubo M, Kimura Y, Odani T, et al. Studies on scutellariae radix. Part II: The antibacterial substance. *Planta Medica* 1981;43:p.194-201.
9. Mahmood N, Pizza C, Aquino R, et al. Inhibition of HIV infection by flavonoids. *Antivir Res* 1993; 22:p.189-99.
10. Ono K, Nakane H, Fukushima M, et al. Differential inhibitory effects of various flavonoids on the activities of reverse transcriptase and cellular DNA and RNA polymerases. *Eur J Biochem* 1990;190:p.469-79.
11. Razina TG, Udintsev SN, Tiutrin II, et al. The role of thrombocyte aggregation function in the mechanism of the antimetastatic action of an extract of Baikal skullcap. *Vopr Onkologii* 1989; 35:p.331-5.
12. Wong, BC, Wang WP, Cho CH, Fan XM Lin MC, Kung HF, et al. 12-Lipoxygenase inhibition induced apoptosis in human gastric cancer cells. *Carcinogenesis* 2001 Sep; vol 22(9):p.1349-54.
13. Timñar J, Rñasño E, Djome B, Li L, Grignon D, Nie D, Honn KV, et al. Expression, subcellular localization and putative function of platelet-type 12-lipoxygenase in human prostate cancer cell lines of different metastatic potential. *Int J Cancer* 2000 Jul 1; vol 87(1):p.37-43.
14. Chang WH, Chen CH, Lu FJ. Different effects of baicalein, baicalin and wogonin on mitochondrial function, glutathione content and cell cycle progression in human hepatoma cell lines. *Planta Med* 2002 Feb;vol 68(2): p.128-32.
15. Ching-Hsein Chen, Huang Lynn LH, Chao-Cheng Huang, Chun-Ching Lin, Fung-Jou Lu, Yashang Lee. Baicalein, a novel apoptotic agent for hepatoma cell lines: a potential medicine for hepatoma. *Nutrition & Cancer* 2000;vol 38 (2):p.287-96.
16. Hsu SL, Hsieh YC, Hsieh WC, Chou CJ. Baicalein induces a dual growth arrest by modulating multiple cell cycle regulatory molecules. *Eur J Pharmacol* 2001 Aug 17; vol 425(3):p.165-71.

17. Chan FL, Choi HL, Chen ZY, Chan PS, Huang Y. Induction of apoptosis in prostate cancer cell lines by a flavonoid, baicalin. *Cancer Lett* 2000 Nov 28; vol 160(2):p.219-28.
18. Liu ZL, Tanaka S, Horigome H, Hirono T, Oka K. Induction of apoptosis in human lung fibroblasts and peripheral lymphocytes in vitro by Shosaiko-to derived phenolic metabolites. *Biol Phar Bull* 2002 Jan; vol 25(1):p.37-41.
19. Ikemoto S, Suygimura K, Yoshida N, Yasumoto R, Wada S, Yamamoto K, Kishimoto T. Antitumor effects of Scutellariae radix and its components baicalein, baicalin, and wogonin on bladder cancer cell lines. *Urology* 2000 Jun; vol 55(6):p.951-5.
20. Hara H, Sakamoto T, Ohtaka H, et al. Effects of baicalein and alpha-tocopherol on lipid peroxidation, free radical scavenging activity and 12-O-tetradecanoylphorbol acetate-induced ear edema. *Eur J Pharmacol* 1992;221:p.193-198.
21. Ueng YF, Shyu CC, Liu TY, Oda Y, Lin YL, Liao JF, et al. Protective effects of baicalein and wogonin against benzo[a]pyrene- and aflatoxin B(1) -induced genotoxicities. *Biochem Pharmacol* 2001 Dec 15;vol 62(12):p.1653-60.
22. Higashitanai A, Tabata S, Hayashi T, Hotta Y. Plant saponins can affect DNA recombination in cultured mammalian cells. *Cell Struct Funct* 1989;14:p.617-24.
23. Hiipakka RA, Zhang HZ, Dai W, Dai Q, Liao S. Structure-activity relationships for inhibition of human 5-alpha-reductases by polyphenols. *Biochem Pharmacol* 2002 Mar 15; vol 63(6):p.1165-76.
24. Small EJ, Frohlich MW, Bok R, Shinohara K, Grossfeld G, Rozenblat Z, et al. Prospective trial of the herbal supplement PC-SPES in patients with progressive prostate cancer. *J Clin Oncol* 2000 Nov 1;vol 18(21): p.3595-603.
25. de la Taille A, Hayek OR, Burchardt M, Burchardt T, Katz AE. Role of herbal compounds (PC-SPES) in hormone-refractory prostate cancer: two case reports. *Journal of Alternative and Complementary Medicine* 2000 Oct; 6(5):p.449-51.
26. Porterfield H. UsToo PC-SPES surveys: review of studies and update of previous survey results. *Mol Urol*, 2000 Fall; vol 4(3):p.289-91;discussion 293.
27. Oh WK, George DJ; Hackmann K, Manola J, Kantoff PW. Activity of the herbal combination, PC-SPES, in the treatment of patients with androgen-independent prostate cancer. *Urology* 2001 Jan; vol 57(1):p.122-6.g
28. Tsai Wayne Y, Chang Wen-Huei, Chen Ching-Hsein, Lu Fung-Jou. Enhancing effect of patented whey protein isolate (Immunocal) on cytotoxicity of an anticancer drug. *Nutrition and Cancer* 2000;38(2):p.200-8.
29. Chen Ching-Hsein, Huang Lynn LH, Huang Chao-Cheng, Lin Chun-Ching, Lee Yashang, Lu Fung-Jou. Baicalein, a novel apoptotic agent for hepatoma cell lines: a potential medicine for hepatoma. *Nutrition and Cancer* 2000;38(2):p.287-95.
30. So FV, Guthrie N, Chambers AF, Moussa M, Carroll KK. Inhibition of human breast cancer cell proliferation and delay of mammary tumorigenesis by flavonoids and citrus juices. *Nutr Cancer* 1996;26:p.167-181.
31. So FV, Guthrie N, Chambers AF, Carroll KK. Inhibition of proliferation of estrogen receptor-positive MCG-7 human breast cancer cells by flavonoids in the presence and absence of excess estrogen. *Cancer Lett* 1997;112:p.127-33.
32. Ahn HC, Lee SY, Kim JW, Son WS, Shin CG, lee BJ. Binding aspects of baicalein to HIV-1 integrase. *Mol Cells* 2001 Aug 31;vol12(1):p.127-30.
33. Wu JA, Attele AS, Zhang L, Yuan C. Anti-HIV activity of medicinal herbs: usage and potential development. *American Journal of Chinese Medicine* 2001;29(1):p.69-81.
34. Nagai HK, Osuga A, Koda A. Inhibition of hypersensitivity reactions by soluble derivatives of baicalein. *Jap J Pharmacol* 1975; 25:p.763-72.
35. Brigham KL. Mediators of the inflammatory process: prostanoids. *Handbook of Inflammation* 1989; vol 6:Mediators of the Inflammatory Process, Elsevier Science Publishers.
36. Alcaez MJ, Ferrandiz ML. Modification of arachidonic metabolism by flavonoids. *J Ethnopharmacol* 1987;21:p.209-29.
37. Kubo M, Matsuda H, Tanaka M, et al. Studies on scutellariae radix VII. Anti-arthritic and

anti-inflammatory actions of methanolic extract and flavonoid components from scutellaria radix. *Chem Pharm Bull* 1984;32:p.2724-9.

38. Lebeau A, Esclaire F, Rostène W, Pélaprat D. Baicalein protects cortical neurons from beta-amyloid (25-35) induced toxicity. *Neuroreport* 2001 Jul 20;vol 12(10):p.2199-202.
39. Gao Z, Huang K, Xu H. Protective effects of flavonoids in the roots of scutellaria baicalensis georgi against hydrogen peroxide-induced oxidative stress in HS-SY5Y cells. *Pharmacol Res* 2001 Feb;vol 43(2):p.173-8.
40. Shieh DE, Liu LT, Lin CC. Antioxidant and free radical scavenging effects of baicalein, baicalin and wogonin. *Anticancer Res* 2000 Sep-Oct; vol 20(5A):p.2861-5.
41. Tierra LL. *The Herbs of Life; Health and Healing Using Western and Chinese Techniques*. Freedom, CA; The Crossing Press, 1992.
42. Healthnotes Online. 2001 Healthnotes Inc. www.healthnotes.com : Chinese skullcap.

James Meschino, DC, MS
Toronto, Ontario
Canada
www.renaissance.com

OCTOBER 2003